

# THE CONNECTION BETWEEN JETS, ACCRETION DISKS, AND BLACK HOLE MASS IN BLAZARS

JIAN-MIN WANG<sup>1</sup>, BIN LUO<sup>1</sup> AND LUIS C. HO<sup>2</sup>

Received 2004 August 17; accepted 2004 September 22; published October 4

## ABSTRACT

The properties of relativistic radio jets are thought to be closely connected with the properties of accretion disks in active galactic nuclei. We explore this issue using a sample of 35 blazars with very-long baseline observations, for which we can estimate the kinetic powers of their relativistic jets, the bolometric luminosity of their accretion disks, and the masses of their black holes. Contrary to previous claims, we find that the jet kinetic power is significantly correlated with the disk luminosity. This supports the notion that the disk is somehow coupled to the jet, even on parsec scales. Moreover, we show that the correlation improves by including the black hole mass as a second parameter. The dominance of the jet, parameterized as the ratio of the jet kinetic power to the disk luminosity, is largely controlled by and is inversely correlated with the Eddington ratio of the accretion disk. This empirical relation should serve as a useful guide for theoretical models for jet formation.

*Subject headings:* accretion, accretion disks — black hole physics — galaxies: active — (galaxies:) BL Lacertae  
 objects: general — galaxies: nuclei — (galaxies:) quasars: general

## 1. INTRODUCTION

Many of the main properties of blazars can be explained by relativistic jets (see review by Urry & Padovani 1995); however, jet formation remains one of the unsolved fundamental problems in astrophysics (e.g., Meier, Koide, & Uchida 2001). Three scenarios have been suggested for the origin of jets: (1) from the vicinity of the black hole (BH) event horizon via extraction of the spin energy of the BH (Blandford & Znajek 1977); (2) from a wind from a magnetized accretion disk (Blandford & Payne 1982); and (3) from an advection-dominated accretion flow (ADAF; Narayan & Yi 1994), either dominated by outflows (Blandford & Begelman 1999) or a jet (Falcke, Körding, & Markoff 2003). Though there are still some unresolved issues with each of these models, it is clear that an accretion disk is involved in all of them. It is expected that there is a connection between the relativistic jet and the accretion disk.

The jet-disk connection has been extensively explored by many authors and in different ways. Since emission from the relativistic jets is highly beamed, one needs to use isotropic radiation to investigate the connection. On large scales ( $\sim$ kpc–Mpc), many authors have tested the correlation between different emission-line luminosity and radio luminosity (Stockton & MacKenty 1987; Baum & Heckman 1989; Rawlings & Saunders 1991; Falcke, Malkan, & Biermann 1995; Xu, Livio, & Baum 1999; Cao & Jiang 2001; Wang, Ho, & Staubert 2003). Indeed, a strong link between the jet and the disk has been found, although we should note that it is indirect because the extended radio emission and the line emission have quite different lifetimes. On  $\sim$ pc scales, radio data from very long-baseline interferometry (VLBI) have been used to calculate the kinetic luminosity to test the jet-disk connection (Celotti, Padovani, & Ghisellini 1997; hereafter CPG). CPG found no correlation between the broad-line region (BLR) luminosity ( $L_{\text{BLR}}$ ) and the kinetic luminosity ( $L_{\text{kin}}$ ), although there are some suggestive hints of a jet-disk connection. Maraschi & Tavecchio (2003; hereafter MT03; see also Maraschi 2001) investigated the problem using a sample of blazars for which the jet power could be

estimated from modeling of their spectral energy distribution. They found that the jet power is linearly proportional to the disk power, with flat-spectrum radio quasars radiating at a greater efficiency than BL Lac objects. In view of the importance of this issue, it certainly merits reexamination.

Jet formation may depend on the state of the accretion disk. Marscher et al. (2002) found that the radio galaxy 3C 120, as in microquasars, ejected superluminous radio blobs following a dip in the X-ray emission. It is essential to probe the accretion rates of the disks to test the jet-disk connection. Wang et al. (2002, 2003, 2004) showed that most BL Lac objects have optically thin ADAFs, while most radio-loud quasars have standard accretion disk. It is interesting to note that the degree of radio-loudness in low-luminosity active galactic nuclei (AGNs) is determined by the Eddington ratio (Ho 2002), and BH accreting systems obey a “fundamental plane” defined by their radio emission, hard X-ray emission, and BH mass (Merloni, Heinz, & Di Matteo 2003). The highly anisotropic emission from the jet depends on the dissipation of the jet kinetic energy, and the radiation efficiency may be controlled by both the state of the central engine and the environment of the jet (Celotti & Fabian 1993; Wang, Staubert, & Ho 2002). However, how the kinetic luminosity, a more fundamental parameter of the relativistic jet, depends on the properties of the accretion disk is not yet known.

In this paper, we tackle this issue with the sample of blazars from CPG. We show that in blazars there is a well-defined correlation between kinetic luminosity, BLR luminosity, which we use as a surrogate to represent the bolometric luminosity of the accretion disk, and the BH mass. Moreover, the degree of jet-dominance diminishes with increasing Eddington ratio.

## 2. THE SAMPLE

For the purposes of the present paper, we need the kinetic luminosity, bolometric luminosity, and the BH mass. According to CPG, the kinetic luminosity is given by  $L_{\text{kin}} = f\pi R_{\text{VLBI}}^2 n_e m_e c^2 \Gamma^2 \beta c$ , where  $R_{\text{VLBI}}$  is the size of the radio jet,  $\beta$  is the jet velocity in units of the light speed ( $c$ ),  $n_e$  is the number density of relativistic electrons,  $\Gamma$  is the Lorentz factor of the jet, and  $m_e$  is the electron mass.  $R_{\text{VLBI}}$  is an observable pa-

<sup>1</sup> Laboratory for High Energy Astrophysics, Institute of High Energy Physics, Chinese Academy of Science, Beijing 100039, P. R. China.

<sup>2</sup> The Observatories of the Carnegie Institution of Washington, 813 Santa Barbara Street, Pasadena, CA 91101-1292, USA

TABLE 1: THE SAMPLE<sup>a</sup>

| Name                  | Type | $z$   | $\mathcal{D}^b$ | $\log L_{\text{kin}}$<br>( $\text{erg s}^{-1}$ ) | $\log L_{\text{BLR}}$<br>( $\text{erg s}^{-1}$ ) | Line       | Ref.  | $\log M_{\text{BH}}$<br>( $M_{\odot}$ ) | $M_{\text{BH}}$ Estimator                | $M_{\text{B}}$<br>(mag) | Ref.  |
|-----------------------|------|-------|-----------------|--------------------------------------------------|--------------------------------------------------|------------|-------|-----------------------------------------|------------------------------------------|-------------------------|-------|
| (1)                   | (2)  | (3)   | (4)             | (5)                                              | (6)                                              | (7)        | (8)   | (9)                                     | (10)                                     | (11)                    | (12)  |
| 0336–019              | HPQ  | 0.852 | 15.6            | 46.21                                            | 44.97                                            | H $\beta$  | JB91  | 8.95                                    | FWHM(H $\beta$ )=4876 km s <sup>-1</sup> | -24.24                  | WU02  |
| 0420–014              | HPQ  | 0.915 | 16.8            | 47.07                                            | 45.12                                            | Mg II      | SF97  | 8.95                                    | FWHM(H $\beta$ )=3000 km s <sup>-1</sup> | -25.84                  | WU02  |
| 0521–365              | HPQ  | 0.055 | 6.17            | 46.18                                            | 42.00                                            | H $\beta$  | SF97  | 8.65                                    | $\sigma$ =269 km s <sup>-1</sup>         | ...                     | WU02  |
| 1034–293              | HPQ  | 0.312 | 5.4             | 45.81                                            | 42.80                                            | H $\beta$  | SF97  | 8.66                                    | FWHM(H $\beta$ )=4116 km s <sup>-1</sup> | -23.69                  | WU02  |
| 1253–055              | HPQ  | 0.538 | 18              | 46.11                                            | 44.23                                            | Mg II      | SF97  | 8.48                                    | FWHM(H $\beta$ )=3100 km s <sup>-1</sup> | -23.93                  | WU02  |
| 1335–127              | HPQ  | 0.541 | 11.9            | 46.50                                            | 44.15                                            | Mg II      | SF97  | 8.42                                    | FWHM(Mg II)=4602 km s <sup>-1</sup>      | -22.94                  | SKF93 |
| 1510–089              | HPQ  | 0.361 | 14.5            | 45.48                                            | 44.59                                            | H $\beta$  | SM87  | 8.62                                    | FWHM(H $\beta$ )=3180 km s <sup>-1</sup> | -24.41                  | WU02  |
| 1641+399              | HPQ  | 0.595 | 5.3             | 44.80                                            | 45.01                                            | H $\beta$  | JB91  | 9.45                                    | FWHM(H $\beta$ )=5140 km s <sup>-1</sup> | -25.94                  | WU02  |
| 1741–038              | HPQ  | 1.054 | 4.3             | 46.79                                            | 44.44                                            | Mg II      | C97   | 9.29                                    | FWHM(Mg II)=11574 km s <sup>-1</sup>     | -23.32                  | F04   |
| 1921–293              | HPQ  | 0.352 | 18.2            | 46.86                                            | 44.02                                            | Mg II      | SF97  | 9.01                                    | FWHM(H $\beta$ )=9134 km s <sup>-1</sup> | -22.40                  | JB91  |
| 2223–052              | HPQ  | 1.404 | 20.9            | 46.46                                            | 45.28                                            | Mg II      | SF97  | 8.81                                    | FWHM(C IV)=3449 km s <sup>-1</sup>       | -25.47                  | W95   |
| 2251+158              | HPQ  | 0.859 | 6               | 46.19                                            | 45.75                                            | H $\beta$  | JB91  | 9.10                                    | FWHM(H $\beta$ )=2800 km s <sup>-1</sup> | -26.64                  | WU02  |
| 2345–167              | HPQ  | 0.576 | 10.8            | 45.84                                            | 44.32                                            | H $\beta$  | JB91  | 8.76                                    | FWHM(H $\beta$ )=4999 km s <sup>-1</sup> | -23.44                  | WU02  |
| 0333+321              | LPQ  | 1.258 | 16.6            | 47.47                                            | 46.30                                            | Mg II      | C97   | 10.11                                   | FWHM(Mg II)=14700 km s <sup>-1</sup>     | -26.59                  | SS91  |
| 0923+392              | LPQ  | 0.699 | 11.6            | 46.03                                            | 45.23                                            | H $\beta$  | O84   | 9.40                                    | FWHM(H $\beta$ )=7200 km s <sup>-1</sup> | -24.65                  | WU02  |
| 1226+023              | LPQ  | 0.158 | 6               | 46.09                                            | 45.42                                            | H $\beta$  | SM87  | 9.21                                    | $M_{\text{B}}(\text{host}) = -22.11$ mag | ...                     | MC04  |
| 1928+738              | LPQ  | 0.302 | 4.4             | 45.53                                            | 44.61                                            | H $\beta$  | M96   | 8.72                                    | FWHM(H $\beta$ )=3360 km s <sup>-1</sup> | -24.58                  | WU02  |
| 1954+513              | LPQ  | 1.220 | 6.17            | 47.24                                            | 44.86                                            | Mg II      | C97   | 9.94                                    | FWHM(Mg II)=15241 km s <sup>-1</sup>     | -25.49                  | L96   |
| 2134+004              | LPQ  | 1.936 | 34.6            | 46.00                                            | 44.75                                            | Mg II      | C92   | 8.91                                    | FWHM(Mg II)=2800 km s <sup>-1</sup>      | -27.87                  | C92   |
| 2351+456              | LPQ  | 2.000 | 6.17            | 46.72                                            | 44.76                                            | Mg II      | SK93  | 8.80                                    | FWHM(Mg II)=5118 km s <sup>-1</sup>      | -24.47                  | F04   |
| 0906+430              | LDQ  | 0.670 | 43.4            | 46.66                                            | 44.67                                            | Mg II      | C97   | 8.95                                    | $M_{\text{B}}(\text{host}) = -21.58$ mag | ...                     | MC04  |
| 1222+216              | LDQ  | 0.435 | 1.3             | 45.67                                            | 44.67                                            | H $\beta$  | C97   | 8.17                                    | FWHM(H $\beta$ )=2197 km s <sup>-1</sup> | -23.92                  | F04   |
| 1317+520              | LDQ  | 1.060 | 1.23            | 48.62                                            | 46.01                                            | Mg II      | C97   | 10.05                                   | FWHM(Mg II)=13900 km s <sup>-1</sup>     | -26.54                  | SS91  |
| 1721+343              | LDQ  | 0.206 | 1.23            | 46.97                                            | 44.67                                            | H $\beta$  | R84   | 8.50                                    | FWHM(H $\beta$ )=2880 km s <sup>-1</sup> | -24.25                  | WU02  |
| 1845+797              | LDQ  | 0.057 | 1.23            | 46.31                                            | 43.54                                            | H $\beta$  | R84   | 8.55                                    | FWHM(H $\beta$ )=8850 km s <sup>-1</sup> | -20.76                  | L96   |
| 0235+164              | BL   | 0.940 | 6.5             | 45.69                                            | 43.81                                            | Mg II      | C97   | >10.22                                  | $M_{\text{R}}(\text{host}) > -26.44$ mag | ...                     | U00   |
| 0537–441              | BL   | 0.896 | 11.6            | 46.44                                            | 44.61                                            | Mg II      | SF97  | 8.74                                    | FWHM(Mg II)=3200 km s <sup>-1</sup>      | -26.34                  | W86   |
| 0851+202              | BL   | 0.306 | 8.8             | 46.37                                            | 43.44                                            | Mg II      | SFK93 | 8.79                                    | FWHM(H $\beta$ )=3443 km s <sup>-1</sup> | -24.78                  | SFK89 |
| 1101+384 <sup>c</sup> | BL   | 0.031 | 3.09            | 45.45                                            | 41.34                                            | H $\alpha$ | C97   | 8.29                                    | $\sigma$ =219 km s <sup>-1</sup>         | ...                     | WU02  |
| 1308+326              | BL   | 0.996 | 6.8             | 45.93                                            | 44.63                                            | Mg II      | SK93  | 9.24                                    | FWHM(Mg II)=4016 km s <sup>-1</sup>      | -27.93                  | SK93  |
| 1538+149              | BL   | 0.605 | 1.3             | 44.58                                            | 43.05                                            | Mg II      | SK93  | 8.94                                    | $M_{\text{R}}(\text{host}) = -23.88$ mag | ...                     | U00   |
| 1652+398 <sup>d</sup> | BL   | 0.034 | 1.5             | 45.36                                            | 41.27                                            | H $\alpha$ | C97   | 9.21                                    | $\sigma$ =372 km s <sup>-1</sup>         | ...                     | WU02  |
| 1803+784              | BL   | 0.684 | 8.5             | 45.74                                            | 44.28                                            | Mg II      | SK93  | 8.57                                    | FWHM(Mg II)=2451 km s <sup>-1</sup>      | -26.66                  | L96   |
| 1807+698              | BL   | 0.051 | 3.09            | 45.11                                            | 41.34                                            | H $\beta$  | C97   | 8.52                                    | $M_{\text{R}}(\text{host}) = -23.04$ mag | ...                     | F03   |
| 2200+420              | BL   | 0.069 | 4.4             | 45.13                                            | 42.52                                            | H $\beta$  | V95   | 8.35                                    | $M_{\text{R}}(\text{host}) = -22.69$ mag | ...                     | F03   |

REFERENCES/NOTES:—C92: Corbin 1992; CPG: Celotti et al. 1997; F03: Falomo, Carangelo, & Treves 2003; L96: Lawrence et al. 1996; M96: Marziani et al. 1996; MC04: Marchesini & Celotti 2004; O84: Oke, Shields, & Korycansky 1984; SF97: Scarpa & Falomo 1997; SFK89: Stickel, Fried, & Kühr 1989; SKF93: Stickel, Kühr, & Fried 1993; SFK93: Stickel, Fried, & Kühr 1993; SK93: Stickel & Kühr 1993; SM:87 Stockton & MacKenty 1987; SS91: Steidel & Sargent 1991; U00: Urry et al. 2000; V95: Vermeulen et al. 1995; W86: Wilkes 1986; W95: Wills et al. 1995; WU02: Woo & Urry 2002.

HPQ: core-dominated high-polarization quasars; LPQ: core-dominated low-polarization quasars; LDQ: lobe-dominated quasars; BL: BL Lacs.

<sup>a</sup>All the data are scaled to luminosity distances calculated assuming  $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ,  $\Omega_m = 0.3$ , and  $\Omega_\Lambda = 0.7$ . <sup>b</sup>Mrk 421 has line flux for H $\alpha$  instead of H $\beta$  or Mg II. H $\beta$  or Mg II lines. The upper limit on its core size given in Ghisellini et al. (1993) gives only an upper limit on  $\log L_{\text{kin}}$ . The value of  $\log L_{\text{kin}}$  listed is based on the new VLBI observations of Piner et al. (1999). <sup>d</sup>Mrk 501 has line flux for H $\alpha$  instead of H $\beta$  or Mg II.

parameter, while  $n_e$  and  $\Gamma$  can be obtained by fitting the observed spectrum. However, the largest source of uncertainty for the kinetic luminosity lies in the factor  $f$ , which describes the particle composition of the jet. For a jet composed of a proton-electron plasma,  $f$  has a value of  $\sim 1840$ , while for a pair plasma  $f = 1$ . As in CPG, we take  $f = 1$ . In the present paper, we use a subset of the sample of CPG, which provides  $L_{\text{BLR}}$  and  $L_{\text{kin}}$ . Our subset is limited by the availability of the following data: (1)  $L_{\text{kin}}$ , (2) BH mass estimate, and (3) flux for the H $\beta$   $\lambda 4861$  or Mg II  $\lambda 2800$  line, which, as described below, we use to estimate  $L_{\text{BLR}}$ . We omit those sources with an upper limit on  $L_{\text{kin}}$  because they only have an upper limit on  $R_{\text{VLBI}}$ . Ghisellini et al. (1993) give the Doppler factors of the sample, estimated from the classical limits of synchrotron self-Compton emission. In this study, we estimate BH masses using one of three methods, depending on the availability of the necessary data: (1) the relation between stellar velocity dispersion and BH mass, as fitted by Tremaine et al. (2002); (2) the linewidth-luminosity-mass scaling relation, which is calibrated against local AGNs studied using reverberation mapping (Kaspi et al. 2000); and (3) the correlation between host galaxy luminosity and BH mass,

as given by McLure & Dunlop (2001). When the linewidth (FWHM) of H $\beta$  is not available, we use the FWHM of C IV or Mg II as suggested by Vestergaard (2002) and McLure & Jarvis (2002), respectively, to estimate the BH mass. Our paper analyzes 35 blazars (Table 1) from the sample of CPG. When the BH mass is estimated from the linewidth-luminosity-mass relation, we use an extrapolation of the continuum from the magnitudes given in Véron-Cetty & Véron (2003), assuming a spectrum of the form  $F_\nu \propto \nu^{-0.5}$ .

The uncertainty in the  $M_{\text{BH}} - \sigma$  relation is small,  $\lesssim 0.21$  dex (Tremaine et al. 2002). On the other hand, the uncertainty on the zeropoint of the linewidth-luminosity-mass relation is approximately 0.5 dex (Gebhardt et al. 2000; Ferrarese et al. 2001). We note, however, that this method may systematically overestimate the BH masses in radio-loud quasars if the optical-ultraviolet continuum in these objects is significantly beamed. Unfortunately, it is difficult to estimate the magnitude of this effect, since the amount of beaming is wavelength dependent (e.g., Bregman 1990), and we have no good constraints for the optical-ultraviolet region. According to McLure & Dunlop (2001), the  $M_{\text{BH}} - M_{\text{R}}$  correlation for quasar host galaxies




FIG. 1.— Correlation between jet kinetic luminosity and (a)  $L_{\text{BLR}}$  and (b)  $L_{\text{BLR}}$  and  $L_{\text{Edd}}$  for BL Lac objects (stars), lobe-dominated quasars (open circles), core-dominated, low-polarization quasars (squares), and core-dominated, high-polarization quasars (filled circles). The uncertainties on  $L_{\text{BLR}}$  are based on the standard deviation of  $\text{Mg II}/\text{Ly}\alpha$  and  $\text{H}\beta/\text{Ly}\alpha$  in the composite quasar spectrum of Francis et al. (1991), while the uncertainties on  $L_{\text{kin}}$  are assumed to be 0.5 dex for all sources. The error bars for the abscissa in (b) take into account the uncertainties in  $L_{\text{BLR}}$  and  $M_{\text{BH}}$  (see §2).

has an uncertainty of 0.6 dex. Given the intrinsic uncertainty of the different BH mass estimators and the heterogeneity of our sample, we estimate that the individual BH masses may have an uncertainty as large as  $\sim 1$  dex.

### 3. EMPIRICAL CORRELATIONS

The BLR luminosities given in CPG were derived by scaling several strong emission lines to the quasar template spectrum of Francis et al. (1991), using  $\text{Ly}\alpha$  as a reference. With this approach, CPG find no significant correlation between  $L_{\text{BLR}}$  and  $L_{\text{kin}}$ . We note, however, that some of the key emission lines used in their method have a very large scatter in their ratio with respect to  $\text{Ly}\alpha$ . For example,  $\text{C IV } \lambda 1550/\text{Ly}\alpha = 63 \pm 41$  and  $\text{Mg II } \lambda 2800/\text{Ly}\alpha = 34 \pm 20$ . The variation in the strength of C IV and Mg II relative to  $\text{Ly}\alpha$  stems from genuine object-to-object variations in the detailed physical conditions of the BLR in quasars. Since this scatter propagates directly into an error in  $L_{\text{BLR}}$ , it is possible that this is responsible for the apparent lack of a correlation between  $L_{\text{BLR}}$  and  $L_{\text{kin}}$ . On the other hand, the ratio  $\text{H}\beta/\text{Ly}\alpha$  is largely governed by recombination physics (Netzer 1990) and hence is much better defined. Francis et al. (1991) find  $\text{H}\beta/\text{Ly}\alpha = 22 \pm 4.1$ . Here, we will adopt  $\text{H}\beta$  alone as the fiducial line to estimate  $L_{\text{BLR}}$ ; when  $\text{H}\beta$  is not available, we use Mg II, which has the second smallest scatter relative to  $\text{Ly}\alpha$ . Following the notation of CPG (their Eq. 1),

$$L_{\text{BLR}} = \begin{cases} \frac{\langle L_{\text{BLR}}^* \rangle}{L_{\text{est}}(\text{H}\beta)} = 25.26 L_{\text{H}\beta}, \\ \frac{\langle L_{\text{BLR}}^* \rangle}{L_{\text{est}}(\text{Mg II})} = 16.35 L_{\text{Mg II}}, \end{cases} \quad (1)$$

where  $\langle L_{\text{BLR}}^* \rangle = 555.77$ ,  $L_{\text{est}}(\text{H}\beta) = 22$ , and  $L_{\text{est}}(\text{Mg II}) = 34$ . Table 1 gives the values of  $L_{\text{BLR}}$  estimated from Equation 1.

Figure 1a shows a strong correlation between  $L_{\text{BLR}}$  and  $L_{\text{kin}}$ . We have tested the partial correlation among  $L_{\text{kin}}$ ,  $L_{\text{BLR}}$ , and  $d_L$ , the luminosity distances of the objects, so as to avoid the possible spurious effects caused by their common dependence on the luminosity distance. Using the least square method of multivariate regression, we find that  $\log L_{\text{kin}} = 0.43 \log L_{\text{BLR}} + 0.11 \log d_L$ , with partial coefficient  $r_{d_L} = 0.4$  and  $r_{L_{\text{BLR}}} = 0.62$ . This test shows that the correlation between  $L_{\text{kin}}$  and  $L_{\text{BLR}}$  is intrinsic rather than introduced by the dependence of  $d_L$ .

Since we are exploring the possible dependence of jets on the accretion disks (using  $L_{\text{BLR}}$  as a substitute for the disk bolometric luminosity),  $L_{\text{BLR}}$  should be treated as the independent variable. The least square method gives the following regression fit in Figure 1a (excluding the prominent outlier 3C 345):

$$\log L_{\text{kin}} = (0.37 \pm 0.08) \log L_{\text{BLR}} + (29.78 \pm 3.7), \quad (2)$$

with Pearson's coefficient  $r = 0.62$  and a probability of  $p = 1.0 \times 10^{-4}$  for rejecting the null hypothesis of no correlation. Assuming that the BLR luminosity is due to the reprocessing of the primary continuum radiation from the accretion disk, Equation 2 can be interpreted as evidence for a strong connection between the relativistic jet and the accretion disk. The relation  $L_{\text{kin}} \propto L_{\text{BLR}}^{0.37}$  is much flatter than the linear relation between jet power and disk power found by MT03. It is unclear what the source of discrepancy is. We simply note that our procedure for estimating disk luminosities differs from that of MT03. MT03 obtain their disk luminosities either directly from the optical-UV luminosity of the big blue bump or from the original prescription of CPG.

Despite the strong correlation exhibited in Figure 1a, clearly there is still significant scatter. The scatter around the best-fit line is  $\sigma = 0.37$  dex. Although a significant fraction of this scatter may be due to measurement and/or systematic errors in  $L_{\text{kin}}$  and  $L_{\text{BLR}}$ , it is possible that it indicates that a second parameter is important. A nature choice to investigate is the BH mass, here cast in terms of its Eddington luminosity,  $L_{\text{Edd}}$ . Using the least square method of multivariate regression, we indeed find that  $L_{\text{kin}}$  correlates with both  $L_{\text{BLR}}$  and  $L_{\text{Edd}}$ . Excluding 3C 345 and 0235+164 (its BH mass is an upper limit), we obtain

$$\log L_{\text{kin}} = (0.25 \pm 0.09) \log L_{\text{BLR}} + (0.65 \pm 0.25) \log L_{\text{Edd}} + 5.07 \pm 10.05, \quad (3)$$

with Pearson's coefficient  $r = 0.70$ . The new correlation (Fig. 1b) shows a smaller scatter, which has reduced to  $\sigma = 0.31$  dex. The significance of adding the Eddington luminosity as a new parameter in the correlation (3) can be tested by the  $F$ -test defined by Bevington (1969). We find  $F_\chi = 7.36$  and probability  $p = 1.09 \times 10^{-2}$ . The improvement in the correlation by including  $L_{\text{Edd}}$  as an additional variable suggests that the jet power

depends on both the disk luminosity (or accretion rate) and BH mass. Whether additional variables are required, however, cannot be decided on the basis of the data presented here. The scatter in Figure 1b, though reduced, remains substantial, but the significant uncertainties currently associated with the observables prevent us from assessing how much of the scatter is intrinsic.

Equation 3 can be casted in a different form. Defining the “jet-dominance” factor as  $\mathcal{F}_J = L_{\text{kin}}/L_{\text{bol}}$ , the Eddington ratio as  $\mathcal{E} = L_{\text{bol}}/L_{\text{Edd}}$ , and assuming  $L_{\text{bol}} \approx 10L_{\text{BLR}}$  (Netzer 1990), it follows that

$$\log \mathcal{F}_J = -0.10 \log L_{\text{bol}} - 0.65 \log \mathcal{E} + 4.82. \quad (4)$$

This implies that the relative importance of the jet power compared to the disk luminosity is mainly controlled by, and is inversely dependent on, the Eddington ratio. MT03 reached a qualitatively similar conclusion in their analysis. This relation provides a physical basis for understanding the spectral sequence of blazars (Fossati et al. 1998; Wang et al. 2002, 2004; MT03). This empirical relation may also help guide theoretical models on jet formation.

#### 4. CONCLUSIONS

Using rough estimates of the BH masses and bolometric luminosities for a sample of 35 blazars with VLBI observations,

we find that the kinetic power of the relativistic jet in these objects is strongly coupled both to the accretion disk luminosity and the BH mass. Equivalently, the degree of jet dominance, defined as the ratio of the jet kinetic power to the bolometric luminosity of the disk, is largely controlled by and is inversely correlated with the Eddington ratio of the accretion disk. This empirical relation should provide strong constraints on theoretical models for jet formation and jet-disk coupling.

There is growing evidence in recent years that BH accretion in AGNs provides an important source of energy feedback to their host galaxies and their large-scale environments (see review by Begelman 2004). Relativistic jets from radio-loud sources, in particular, may be particularly effective in heating the intracluster medium in galaxy clusters. The empirical correlations described in this study, if they can be properly calibrated, opens up the possibility of using them as tools to investigate quantitatively accretion-powered feedback in the context of galaxy formation.

This research is supported by a Grant for Distinguished Young Scientists from NSFC and 973 project. The work of LCH is funded by the Carnegie Institution of Washington and by NASA grants from the Space Telescope Science Institute (operated by AURA, Inc., under NASA contract NAS5-26555).

#### REFERENCES

- Baum, S. A., & Heckman, T. M. 1989, *ApJ*, 336, 681  
 Begelman, M. C. 2004, in *Carnegie Observatories Astrophysics Series, Vol. 1: Coevolution of Black Holes and Galaxies*, ed. L. C. Ho (Cambridge: Cambridge Univ. Press), in press  
 Bevington, P. R., 1969, *Data Reduction and Error Analysis for the Physical Sciences*. (New York: McGraw-Hill)  
 Blandford, R. D., & Begelman, M. C. 1999, *MNRAS*, 303, L1  
 Blandford, R. D., & Payne, D. G. 1982, *MNRAS*, 199, 883  
 Blandford, R. D., & Znajek, R. L. 1977, *MNRAS*, 179, 433  
 Bregman, J. N. 1990, *A&A Rev*, 2, 125  
 Cao, X., & Jiang, D. R. 2001, *MNRAS*, 320, 347  
 Celotti, A., & Fabian, A. C. 1993, *MNRAS*, 264, 228  
 Celotti, A., Padovani, P., & Ghisellini, G. 1997, *MNRAS*, 286, 415 (CPG)  
 Corbin, M. R. 1992, *ApJ*, 391, 577  
 Falcke, H., K rding, E., & Markoff, S. 2004, *A&A*, 414, 895  
 Falcke, H., Malkan, M. A., & Biermann, P. L. 1995, *A&A*, 336, 702  
 Falomo, R., Carangelo, N., & Treves, A. 2003, *MNRAS*, 343, 505  
 Ferrarese, L., Pogge, R. W., Peterson, B. M., Merritt, D., Wandel, A., & Joseph, C. L. 2001, *ApJ*, 555, L79  
 Fossati, G., Maraschi, L., Celotti, A., Comastri, A., & Ghisellini, G. 1998, *MNRAS*, 299, 433  
 Francis, P. J., Hewett, P. C., Foltz, C. B., Chaffee, F. H., Weymann, R. J., & Morris, S. L. 1991, *ApJ*, 373, 465  
 Gebhardt, K., et al. 2000, *ApJ*, 543, L5  
 Ghisellini, G., Padovani, P., Celotti, A., & Maraschi, L. 1993, *ApJ*, 407, 65  
 Ho, L. C. 2002, *ApJ*, 564, 120  
 Jackson, N., & Browne, I. W. 1991, *MNRAS*, 250, 414  
 Kaspi, S., Smith, P. S., Netzer, H., Maoz, D., Jannuzi, B. T., & Givon, U. 2000, *ApJ*, 533, 631  
 Lawrence, C. R., Zucker, J. R., Readhead, A. C. S., Unwin, S. C., Pearson, T. J., & Xu, W. 1996, *ApJS*, 107, 541  
 Maraschi, L. 2001, in *The 20th Texas Symposium on Relativistic Astrophysics*, ed. H. Martel & J. C. Wheeler (Melville: AIP), 409  
 Maraschi, L., & Tavecchio, F. 2003, *ApJ*, 593, 667 (MT03)  
 Marchesini, D., & Celotti, A. 2004, *MNRAS*, in press (astro-ph/0403272)  
 Marscher, A. P., et al. 2002, *Nature*, 417, 625  
 Marziani, P., Sulentic, J. W., Dultzin-Hacyan, D., Calvani, M., & Moles, M. 1996, *ApJ*, 410, 56  
 McLure, R. J., & Dunlop, J. S. 2001, *MNRAS*, 327, 199  
 McLure, R. J., & Jarvis, M. J. 2002, *MNRAS*, 337, 109  
 Meier, D., Koide, S., & Uchida, Y. 2001, *Science*, 291, 84  
 Merloni, A., Heinz, S., & Di Matteo, T. 2003, *MNRAS*, 345, 1057  
 Narayan, R., & Yi, I. 1994, *ApJ*, 428, L13  
 Netzer, H. 1990, in *Active Galactic Nuclei*, ed. R. D. Blandford, H. Netzer, & L. Woltjer (Berlin: Springer), 57  
 Oke, J. B., Shields, G. A., & Korycansky, D. G. 1984, *ApJ*, 277, 64  
 Piner, B. G., Unwin, S. C., Wehrle, A. E., Edwards, P. G., Fey, A. L., & Kingham, K. A. 2000, *ApJ*, 525, 176  
 Rawlings, S., & Saunders, R. 1991, *Nature*, 349, 138  
 Scarpa, R., & Falomo, R. 1997, *A&A*, 325, 109  
 Steidel, C. C., & Sargent, W. L. W. 1991, *ApJ*, 382, 433  
 Stickel, M., Fried, J. W., & K hr, H. 1989, *A&AS*, 80, 103  
 Stickel, M., Fried, J. W., & K hr, H. 1993, *A&AS*, 98, 393  
 Stickel, M., & K hr, H. 1993, *A&AS*, 101, 521  
 Stickel, M., K hr, H., & Fried, J. W. 1993, *A&AS*, 97, 483  
 Stockton, A., & MacKenty, J. W. 1987, *ApJ*, 316, 584  
 Tremaine, S., et al. 2002, *ApJ*, 574, 740  
 Urry, C. M., & Padovani, P. 1995, *PASP*, 107, 803  
 Urry, C. M., Scarpa, R., O’Dowd, M., Falomo, R., Pesce, J. E., & Treves, A. 2000, *ApJ*, 532, 816  
 V ron-Cetty, M.-P., & V ron, P. 2003, *A&A*, 412, 399  
 Vestergaard, M. 2002, *ApJ*, 571, 733  
 Vermeulen, R. C., Ogle, P. M., Tran, H. D., Browne, I. W. A., Cohen, M. H., Readhead, A. C. S., Taylor, G. B., & Goodrich, R. W. 1995, *ApJ*, 452, L5  
 Wang, J.-M., Ho, L. C., & Staubert, R. 2003, *A&A*, 409, 887  
 Wang, J.-M., Ho, L. C., & Staubert, R. 2004, in *Carnegie Observatories Astrophysics Series, Vol. 1: Coevolution of Black Holes and Galaxies*, ed. L. C. Ho (Pasadena: Carnegie Observatories, <http://www.ociw.edu/ociw/symposia/series/symposium1/proceedings.html>)  
 Wang, J.-M., Staubert, R., & Ho, L. C. 2002, *ApJ*, 579, 554  
 Wilkes, B. J. 1986, *MNRAS*, 218, 331  
 Wills, B. J., Brotherton, M. S., Fang, D., Steidel, C. C., & Sargent, W. L. W. 1993, *ApJ*, 415, 563  
 Woo, J.-H., & Urry, C. M. 2002, *ApJ*, 579, 530  
 Xu, C., Livio, M., & Baum, S. A. 1999, *AJ*, 118, 1169